Adult/Adolescent AIDS Cases and Rate per 100,000, by Size of Place of Residence* and Region, Reported in 1999, United States#

	Metropolitan area >500,000		Metropolitan area 50,000 - 500,000		Nonmetropolitan area	
Region	Number	<u>Rate</u>	Number	<u>Rate</u>	Number	Rate
Northeast	12,671	38	807	17	411	9
North Central	3,408	12	510	5	376	3
South	13,375	32	2,792	17	2,187	11
West	7,071	21	485	7	295	5



^{*}Based on reported residence at AIDS diagnosis *Cases from Puerto Rico, US Virgin Islands and territories are not included

Adult/Adolescent AIDS Cases and Rate per 100,000 Population, by Size of Place of Residence and Region, Reported in 1999, United States

In each region of the United States, the majority of persons reported with AIDS are from large metropolitan areas, followed by smaller metropolitan areas, and the fewest from nonmetropolitan areas. The South has the largest proportion of cases reported from nonmetropolitan areas, and the highest rate in those places. Although the large metropolitan areas have the most AIDS cases, the smaller metropolitan and nonmetropolitan areas, especially in the South, also share a significant burden of the AIDS epidemic. These places may face different challenges than the larger areas to provide adequate care and services to the affected populations.

